



(21) (A1)	2,100,212
(22)	1993/07/09
(43)	1994/01/10

5,065,9/52

(51) INTL.CL.<sup>5</sup> A61K-037/54; A61K-037/02

(19) (CA) **APPLICATION FOR CANADIAN PATENT** (12)

(54) Use of Complement Inhibitors for the Preparation of a  
Pharmaceutical for the Prophylaxis and Therapy of  
Inflammatory Intestinal and Skin Disorders as Well as  
Purpura

(72) Römisch, Jürgen - Germany (Federal Republic of) ;  
Pâques, Eric-Paul - Germany (Federal Republic of) ;  
Bartlett, Robert - Germany (Federal Republic of) ;  
Dickneite, Gerhard - Germany (Federal Republic of) ;

(71) Behringwerke Aktiengesellschaft - Germany (Federal  
Republic of) ;

(30) (DE) P 42 22 534.5 1992/07/09

(57) 9 Claims

Notice: This application is as filed and may therefore contain an  
incomplete specification.

**Canada**

2100212

**BEHRINGWERKE AKTIENGESELLSCHAFT 92/B 016 - Ma 955**

Abstract of the disclosure

**The use of complement inhibitors for the preparation of a pharmaceutical for the prophylaxis and therapy of inflammatory intestinal and skin disorders as well as purpura**

---

The invention relates to the use of complement inhibitors, especially of C1 inactivator or of factors I or H, for the preparation of a pharmaceutical for the prophylaxis and therapy of chronic inflammatory intestinal disorders, inflammatory skin disorders and purpura.

5     The use of complement inhibitors for the preparation of  
a pharmaceutical for the prophylaxis and therapy of  
inflammatory intestinal and skin disorders as well as  
purpura

10     The invention relates to the use of complement inhibitors, especially of C1 inactivator or of factors I or H, for the preparation of a pharmaceutical for the prophylaxis and therapy of chronic inflammatory intestinal disorders, inflammatory skin disorders and purpura.

15     The complement system is composed of a number of proteins, most of them proteases, which after initial activation lead to the formation of the terminal lysis complex and thus to the destruction of target cells. Such processes are initiated "classically" by formation of immune complexes or contact activation and "alternatively" by exogenous structures such as bacteria and their lipopolysaccharides. The two  
20     activation pathways merge in the production of component C3b which, together with C5-C9, initiates the membrane attack complex. Proteolytic activation of components C3, C4 and C5 leads to release of the anaphylatoxins C3a, C4a and C5a, which have chemotactic effects on inflammatory  
25     cells.

30     The principal physiological regulators of the complement system are the proteins which have inhibitory activity - C1 inactivator, factor I (also called C3b inactivator) and its accelerator factor H. The first-mentioned inhibitor displays its effect at the site of initiation of the "classical" pathway by interaction with the activating protease. By contrast, factor I is a protease whose catalytic action is considerably increased by factor H and which inactivates the C3b molecule as well as the C4b

molecule by partial degradation and thus intervenes in a regulatory/inhibitory manner at the merging of the "classical" and "alternative" pathway.

5      Activation of the complement system has been observed  
during the course of a number of autoimmune diseases,  
including lupus erythematosus and rheumatoid arthritis.  
These activation processes are generally associated with  
a consumption of the factors involved, especially of the  
10      inhibitors. Although an increase in the complement  
factors may often be observed during the course of the  
acute phase reaction, the inhibitory capacity is insuffi-  
cient to control complement activation and the con-  
sequences resulting therefrom. Replacement of these  
regulators or prophylactic administration therefore  
15      appears worthwhile. The therapeutic value of factors I  
and H for glomerulonephritis has been described in  
EP-A-0 222 611.

The etiopathogenesis of chronic inflammatory intestinal  
disorders, especially "Crohn's disease" and "ulcerative  
20      colitis" has not been explained to date. A "primary  
stimulus" initiates an immune process which is followed  
by tissue infiltration of inflammatory cells and finally  
damage to cells and tissues. The clinical signs are  
ulcers, fistulations and frequent hemorrhagic  
25      stools/diarrhoea, which are associated with attacks of  
fever and spasmodic pain.

It has not to date been possible to define for these  
disorders pathognomonic autoantibodies or autoreactive T  
cells which identify such inflammatory processes as  
30      autoimmune disease. Involvement of immune processes which  
may result in complement activation is not ruled out,  
however. It is moreover possible for the resulting  
anaphylatoxins C3a, C4a and C5a to contribute to  
attracting inflammatory cells. Also noteworthy are the  
35      edemas of the intestinal mucosa which occur in the early

phase of the disorder and are typically observed after activation of the complement system.

5 Elevated C5a levels are also measured in skin disorders such as, for example, in pustular dermatoses, dermatitides or psoriasis and unambiguously prove the activation of the complement system and mediate the attraction of inflammatory cells. The latter in turn make a crucial contribution to the severity of the clinical picture.

10 It is also possible in the clinical picture of purpura for, for example, endothelial cells to be subject to edematous changes leading to capillary dilatation. The clinical manifestation of this condition is erythema/edema, for example, at the site of the cutaneous lesion. The term "purpura" is generally understood to mean the  
15 extravasation of "formed elements of the blood" from dermal blood vessels into the skin (for example dermis). In idiopathic thrombocytopenic purpura there is typically a tendency to bleed with hematoma formation (extravasation), which is crucially assisted by a depletion of  
20 blood platelets. An "inflammatory" purpura is frequently associated with vasculitis which may be caused by immune complexes.

Accordingly, chronic inflammatory intestinal disorders as well as purpura (associated with dermal necroses) and  
25 inflammatory skin disorders have the extravasation of fluid (from blood vessels) in common. The formation of edema which frequently results, as well as the attraction and infiltration of inflammatory cells into the inflammatory tissue are processes which are frequently observed  
30 after activation of the complement system.

The animal model known as the "Arthus reaction" is frequently used for investigation of the described processes and for the pharmacological testing of therapeutically active substances (see Example 1).

We have now found that the complement inhibitors C1 inactivator and the factors I and H have an inhibitory effect on the Arthus reaction.

5 It is accordingly possible to use complement inhibitors, especially C1 inactivator and/or factor I and/or factor H-containing solutions, for the prophylaxis and therapy of inflammatory skin disorders such as, for example, pustular dermatoses, dermatitides or psoriasis and intestinal disorders, especially Crohn's disease and  
10 ulcerative colitis, as well as inflammatory purpura (associated with dermal necroses).

The invention relates to the use of complement inhibitors for the preparation of a pharmaceutical for the prophylaxis and therapy of chronic inflammatory intestinal  
15 disorders, inflammatory skin disorders and purpura.

The C1 inactivator and the factors I and H or combinations of these are particularly suitable.

Purified inhibitors, which can be prepared from blood plasma in a manner known to the person skilled in the  
20 art, are preferably used.

Inhibitors expressed by genetic engineering means and purified can also be used for this purpose.

The dosages used on administration by the intravenous (bolus or infusion), intramuscular or subcutaneous route  
25 are as follows:

C1 inactivator : 1-5000 IU/kg body weight (BW) per day, preferably 5-500 IU/kg × day.

Factor I : 0.005-100 mg/kg BW per day, preferably 0.01-50 mg/kg × day.

Factor H : 0.005-100 mg/kg BW per day, preferably 0.01-50 mg/kg x day.

5 The inhibitors can be used separately or in combination. In general, administration of only one of these inhibitors is sufficient.

10 A combination of the inhibitory proteins appears worthwhile especially when the activation of two complement pathways, the classical and the alternative, cannot be ruled out. Although C1 inactivator alone inhibits contact  
15 activation of the classical pathway and thus also the subsequent release of the anaphylatoxins, it has no effect on the alternative pathway. By contrast, production of the terminal lysis complex can be controlled by the use of factors I and/or H. Consumption  
of the complement factors up to the point of merging of the two activation pathways cannot, by contrast, be stopped by use of factors I and H alone.

**Example:**

20 The Arthus reaction, identified as acute, necrotizing, inflammatory lesion of blood vessels, is induced by immunization and subsequent challenge with an exogenous antigen or directly with an antibody directed against the test species. Extravasation, formation of edema and attraction of inflammatory cells characterize the Arthus  
25 reaction. The parameter measured is the extent of the formation of edema in the rear paw of a rat. The better the reduction/prevention of this formation of edema, the more effective is the substance tested as therapeutic agent.

30 The results of the investigation of the complement inhibitors on the Arthus reaction are listed in Table 1. Both C1 inactivator and factor H as well as factor I have an inhibitory effect on the formation of edema after

administration of an i.v. bolus. Factor I significantly prevents swelling on subcutaneous administration too.

Table 1: Effect of C1 inactivator, factor I and factor H on the Arthus reaction in the rat paw

- 5 The inhibition (%) of swelling was determined by comparison with a control group (placebo). A group was treated with prednisolone as positive control. The test substances were administered one hour before challenge (i.v. = intravenous; s.c. = subcutaneous; p.o. = oral),  
10 and the swelling of the paw was measured four hours thereafter (10 animals/group).

	Substance	Dose	Administration	Inhibition (%)
	C1 inactivator	100 IU/kg	i.v.	12
		200 IU/kg	i.v.	38
15	Factor I	0.05 mg/kg	i.v.	43
		0.50 mg/kg	i.v.	55
		5.00 mg/kg	i.v.	65
		10.00 mg/kg	s.c.	59
		20.00 mg/kg	s.c.	57
20	Factor H	1.0 mg/kg	i.v.	35
		10.0 mg/kg	i.v.	44
	Prednisolone	29.0 mg/kg	p.o.	70



THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE  
PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. The use of complement inhibitors for the preparation  
of a composition for the therapy and prophylaxis of  
inflammatory skin disorders and intestinal disorders  
as well as purpura.
2. The use as claimed in claim 1, wherein C1  
inactivator, factor I or factor H or a combination  
of these is used as complement inhibitor.
3. The use as claimed in claim 1, wherein a composition  
for the therapy and prophylaxis of Crohn's disease  
or ulcerative colitis is prepared.
4. The use as claimed in claim 2, wherein a composition  
for the therapy and prophylaxis of Crohn's disease  
or ulcerative colitis is prepared.
5. The use as claimed in claim 1, wherein a composition  
for the therapy and prophylaxis of pustular  
dermatoses, dermatitides or psoriasis is prepared.
6. The use as claimed in claim 2, wherein a composition  
for the therapy and prophylaxis of pustular  
dermatoses, dermatitides or psoriasis is prepared.
7. The use as claimed in claim 1, wherein a composition  
containing 1-5000 IU/kg x day C1 inactivator or  
0.005-100 mg/kg x day factor I or  
0.005-100 mg/kg x day factor H is prepared.
8. The use as claimed in claim 1, wherein a composition  
containing 5-500 IU/kg x day C1 inactivator or  
0.01-50 mg/kg x day factor I or 0.01-50 mg/kg x day  
factor H is prepared.

9. The use as claimed in claim 1, wherein a composition which can be administered intravenously, intramuscularly or subcutaneously is prepared.